


# California Wildlife Habitat Relationships System

California Department of Fish and Game

California Interagency Wildlife Task Group

**The correct name is  
Melozone crissalis**

California Towhee  
*Pipilo crissalis*  
B484



Range maps are based on available occurrence data and professional knowledge. They represent current, but not historic or potential, range. Unless otherwise noted above, maps were originally published in Zeiner, D.C., W.F. Laudenslayer, Jr., K.E. Mayer, and M. White, eds. 1988-1990. California's Wildlife. Vol. I-III. California Depart. of Fish and Game, Sacramento, California. Updates are noted in maps that have been added or edited since original publication.

**California Wildlife Habitat Relationships System**  
**California Department of Fish and Game**  
**California Interagency Wildlife Task Group**

---

CALIFORNIA TOWHEE

*Pipilo crissalis*

Family: EMBERIZIDAE  
B484

Order: PASSERIFORMES

Class: AVES

Written by: D. Dobkin, S. Granholm

Reviewed by: L. Mewaldt

Edited by: R. Duke

#### DISTRIBUTION, ABUNDANCE, AND SEASONALITY

The former brown towhee recently has been split into the California towhee and the canyon towhee, *P. fuscus* (American Ornithologists' Union 1989). The California towhee is a common, characteristic resident of foothills and lowlands in most of cismontane California. Frequents open chaparral and coastal scrub, as well as brushland patches in open riparian, hardwood-conifer, cropland, and urban habitats. Commonly uses edges of dense chaparral and brushy edges of densely wooded habitats. Also occurs in lowest montane habitats of similar structure in southern California, and locally in Siskiyou and western Modoc cos. Local on coastal slope north of southern Humboldt Co., and apparently absent from western San Joaquin Valley (Grinnell and Miller 1944, McCaskie et al. 1979, Garrett and Dunn 1981). The Inyo California towhee, *P. c. eremophilus*, occurs only in the Argus Mountains of southwestern Inyo Co.

#### SPECIFIC HABITAT REQUIREMENTS

**Feeding:** Feeds on seeds, insects, and some fruits. Gleans and scratches in litter, picks seeds and fruits from plants, and rarely flycatches (Davis 1957). Prefers to forage on open ground adjacent to brushy cover. Insects are important in breeding season, often constituting a third of the diet (Martin et al. 1961).

**Cover:** Shrubs in broken chaparral, margins of dense chaparral, willow thickets, and brushy understory of open wooded habitats provide cover.

**Reproduction:** Nest is a bulky cup of thin twigs, forb stems with leaves and flower-heads, and grasses; lined with fine stems, grasses, and hairs. Nest concealed in low, dense foliage of a shrub or tree, or occasionally on ground. Nest usually placed 0.9 to 3.6 m (3-12 ft) above ground, occasionally as high as 10.7 m (35 ft) (Terres 1980).

**Water:** Drinks freely in captivity and in the wild (Bartholomew and Cade 1963), although in drier habitats may meet most water needs from insects and succulent foods (Dawson 1954).

**Pattern:** Breeds and seeks cover in brush and dense thickets and forages in adjacent open areas.

#### SPECIES LIFE HISTORY

**Activity Patterns:** Yearlong, diurnal activity.

**Seasonal Movements/Migration:** Not migratory.

**Home Range:** Marshall (1960) reported a home range of about 6 ha (15 ac) for the closely related canyon towhee in riparian woodland in Arizona. In broadleaf evergreen habitat in

Alameda Co., Cogswell (1973) reported 29 males per 40 ha (100 ac). Gaines (1974a) reported 10-18 males per 40 ha (100 ac) in Central Valley riparian cover. In California chaparral, Yeaton (1974) found 20 pairs per 40 ha (100 ac). In California oak-riparian habitat, Bontrager (1974) reported 39 males per 40 ha (100 ac).

**Territory:** Territory in California averaged 0.4 to 0.8 ha (1-2 ac) and varied up to 2 ha (5 ac) (Bent 1968). Territory held all year but defended mostly in breeding season.

**Reproduction:** Breeding season early April into early August with peak from early May to mid-June. Apparently a monogamous, solitary breeder. Clutch usually contains 3 or 4 eggs, sometimes 5; may raise 2 or 3 broods each yr. Incubation about 11 days, by female. Altricial young tended by both parents and leave nest at about 8 days. First brood driven away when next brood hatches; otherwise may remain 4-6 wk (Harrison 1978).

**Niche:** Has increased numbers and range as logging, farming, and urbanization have increased habitat. Apparently an uncommon cowbird host (Ehrlich et al. 1988).

**Comments:** The tiny, isolated race of Inyo California towhee is classified as Federal Threatened and California Endangered. Population now restricted to southern Argus Mts.; may be fewer than 150 individuals remaining (California Department of Fish and Game 1989).

## REFERENCES

- American Ornithologists' Union. 1989. Thirty-seventh supplement to the American Ornithologists' Union Check-list of North American birds. *Auk* 106:532-538.
- Bartholomew, G. A., and T. J. Cade. 1963. The water economy of land birds. *Auk* 80:504-539.
- Bent, A. C. (O. L. Austin, Jr., ed.). 1968. Life histories of North American cardinals, grosbeaks, buntings, towhees, finches, sparrows, and allies. 3 Parts. U.S. Natl. Mus. Bull. 237. 1889pp.
- Bontrager, D. R. 1974. Sycamore-coast live oak riparian woodland. Pages 1035-1036 in W. T. Van Velzen, ed. Thirty-eighth breeding bird census. *Am. Birds* 28:987-1054.
- California Department of Fish and Game. 1989. 1988 annual report on the status of California's state listed threatened and endangered plants and animals. Sacramento. 129pp.
- Cogswell, H. L. 1973. Broadleaf evergreen forest with shrub-filled openings. Pages 992-993 in W. T. Van Velzen, ed. Thirty-seventh breeding bird census. *Am. Birds* 27:955-1019.
- Davis, J. 1957. Comparative foraging behavior of the spotted and brown towhees. *Auk* 74:129-166.
- Dawson, W. R. 1954. Temperature regulation and water requirements of the brown and Abert towhees, *Pipilo fuscus* and *Pipilo aberti*. *Univ. Calif. Publ. Zool.* 59:81-124.
- Desante, D. F., and D. G. Ainley. 1980. The avifauna of the South Farallon Islands, California. *Studies in Avian Biol.* No. 4. Cooper Ornithol. Soc., Lawrence, KA. 104pp.
- Ehrlich, P. R., D. S. Dobkin, and D. Wheye. 1988. *The birder's handbook*. Simon and Schuster, New York. 785pp.
- Gaines, D. 1974a. A new look at the nesting riparian avifauna of the Sacramento Valley, California. *West. Birds* 5:61-80.
- Garrett, K., and J. Dunn. 1981. *Birds of southern California*. Los Angeles Audubon Soc. 408pp.
- Grinnell, J., and A. H. Miller. 1944. *The distribution of the birds of California*. Pac. Coast Avifauna No. 27. 608pp.
- Harrison, C. 1978. *A field guide to the nests, eggs and nestlings of North American birds*. W. Collins Sons and Co., Cleveland, OH. 416pp.
- Hensley, M. M. 1954. Ecological relations of the breeding bird population of the desert biome in Arizona. *Ecol. Monogr.* 234:185-207.
- Marshall, J. T., Jr. 1960. Interrelations of Abert and brown towhees. *Condor* 62:49-64.
- Martin, A. C., H. S. Zim, and A. L. Nelson. 1961. *American wildlife and plants, a guide to*

- wildlife food habits. Dover Publ., Inc., New York. 500pp.
- McCaskie, G., P. De Benedictis, R. Erickson, and J. Morlan. 1979. Birds of northern California, an annotated field list. 2nd ed. Golden Gate Audubon Soc., Berkeley. 84pp.
- Terres, J. K. 1980. The Audubon Society encyclopedia of North American Birds. A. Knopf, New York. 1100pp.
- Yeaton, R. I. 1974. An ecological analysis of chaparral and pine forest bird communities on Santa Cruz Island and mainland California. *Ecology* 55:959-973.

B484

---

Life history accounts for species in the California Wildlife Habitat Relationships (CWHR) System were originally published in: Zeiner, D.C., W.F.Laudenslayer, Jr., K.E. Mayer, and M. White, eds. 1988-1990. California's Wildlife. Vol. I-III. California Department of Fish and Game, Sacramento, California. Updates are noted in accounts that have been added or edited since original publication.