


California Wildlife Habitat Relationships System

California Department of Fish and Game

California Interagency Wildlife Task Group

Brown Creeper *Certhia americana* B364



Range maps are based on available occurrence data and professional knowledge. They represent current, but not historic or potential, range. Unless otherwise noted above, maps were originally published in Zeiner, D.C., W.F. Laudenslayer, Jr., K.E. Mayer, and M. White, eds. 1988-1990. California's Wildlife. Vol. I-III. California Depart. of Fish and Game, Sacramento, California. Updates are noted in maps that have been added or edited since original publication.

California Wildlife Habitat Relationships System
California Department of Fish and Game
California Interagency Wildlife Task Group

BROWN CREEPER

Certhia americana

Family: CERTHIIDAE
B364

Order: PASSERIFORMES

Class: AVES

Written by: M. Green

Reviewed by: L. Mewaldt, D. Airola

Edited by: R. Duke, D. Winkler

DISTRIBUTION, ABUNDANCE, AND SEASONALITY

A common to uncommon resident in montane habitats throughout the state, and in coastal conifer habitats south to San Luis Obispo Co. There is downslope movement to interior and coastal foothills and lowlands in fall and winter. A rare transient in southern deserts and on Channel Islands in fall and winter. May winter above normal breeding elevation in conifer habitats in the Sierra Nevada (Gaines 1977b). Prefers dense, mature stands of conifers, but also found in hardwood and hardwood-conifer habitats, especially in winter (Grinnell and Miller 1944, Garrett and Dunn 1981).

SPECIFIC HABITAT REQUIREMENTS

Feeding: Gleans insects, their larvae, and other invertebrates, from bark of tree trunks and branches; also probes in crevices in bark. May eat some seeds in eastern U.S. Typically spirals up tree trunk feeding (Bent 1948).

Cover: Dense, mature stands of conifers provide cover. Hardwoods and riparian deciduous trees also used, especially in winter.

Reproduction: Builds nest behind loose bark, or rarely in tree cavity. Typically nests in old-growth incense cedar, coastal redwood, pine or fir, or snag. In Sierra Co., nest tree height averaged 26.5 m (85 ft), range 6-48 m (18-142 ft). Nest height averaged 2.2 m (7 ft), range 0.3-6.5 m (1-20 ft). Average nest tree dbh was 68 cm (26.6 in), range 35-164 cm (14-64 in) (Raphael and White 1984).

Water: No information found.

Pattern: Frequents dense, old-growth coniferous forests. Also uses deciduous riparian trees and other hardwoods, especially in winter.

SPECIES LIFE HISTORY

Activity Patterns: Yearlong, diurnal activity.

Seasonal Movements/Migration: May move downslope in winter. Appearance on lower slopes in southern coastal lowlands, in southern deserts, and on coastal islands may indicate limited latitudinal migration peaking in October (DeSante and Ainley 1980).

Home Range: Probably same as territory, but little information available.

Territory: No information found. Number of pairs, or individuals, per 40 ha (100 ac) reported as: 4-5 pairs in Great Britain (Jones 1972), 1.7 pairs in burned Jeffrey pine-fir stands and 3.6 in unburned stands in the Sierra Nevada (Bock and Lynch 1970), 24 in a Marin Co.

bay-Bishop pine-mixed forest (Stewart 1974).

Reproduction: Monogamous. Pair breeds solitarily from March to mid-August, with peak activity from mid-May to mid-June. Clutch averages 4 eggs; range 3-5. Incubation 14-17 days; by female at night. Altricial young fledge in 15-16 days (Harrison 1978, Ehrlich et al. 1988).

Niche: Subject to predation from snakes, accipiters and small mammals.

REFERENCES

- Bent, A. C. 1948. Life histories of North American nuthatches, wrens, thrashers, and their allies. U.S. Natl. Mus. Bull. 195. 475pp.
- Bock, C. E., and J. F. Lynch. 1970. Breeding bird populations of burned and unburned conifer forests in the Sierra Nevada. *Condor* 72:182-189.
- DeSante, D. F., and D. G. Ainley. 1980. The avifauna of the South Farallon Islands, California. *Studies in Avian Biol. No. 4.* Cooper Ornithol. Soc., Lawrence, KA. 104pp.
- Ehrlich, P. R., D. S. Dobkin, and D. Wheye. 1988. *The birder's handbook.* Simon and Schuster, New York. 785pp.
- Gaines, D. 1977b. *Birds of the Yosemite Sierra.* California Syllabus, Oakland. 153pp.
- Garrett, K., and J. Dunn. 1981. *Birds of southern California.* Los Angeles Audubon Soc. 408pp.
- Grinnell, J., and A. H. Miller. 1944. The distribution of the birds of California. *Pac. Coast Avifauna No. 27.* 608pp.
- Harrison, C. 1978. *A field guide to the nests, eggs and nestlings of north American birds.* W. Collins Sons and Co., Cleveland, OH. 416pp.
- Jones, P. H. 1972. Succession in breeding bird populations of sample Welsh oakwoods. *Brit. Birds* 65:291-299.
- McCaskie, G., P. De Benedictis, R. Erickson, and J. Morlan. 1979. *Birds of northern California, an annotated field list.* 2nd ed. Golden Gate Audubon Soc., Berkeley. 84pp.
- Raphael, M. G., and M. White. 1984. Use of snags by cavity-nesting birds in the Sierra Nevada. *Wild. Monogr. No. 86.* 66pp.
- Stewart, R. M. 1974. California bay-Bishop pine-mixed forest. Page 1037-1038 in W. T. Van Velzen, ed. *Thirty-eighth breeding bird census.* *Am. Birds* 28:987-1054.

B364

Life history accounts for species in the California Wildlife Habitat Relationships (CWHR) System were originally published in: Zeiner, D.C., W.F.Laudenslayer, Jr., K.E. Mayer, and M. White, eds. 1988-1990. *California's Wildlife.* Vol. I-III. California Department of Fish and Game, Sacramento, California. Updates are noted in accounts that have been added or edited since original publication.