


California Wildlife Habitat Relationships System

California Department of Fish and Game

California Interagency Wildlife Task Group

Chipping Sparrow *Spizella passerina* B489



Range maps are based on available occurrence data and professional knowledge. They represent current, but not historic or potential, range. Unless otherwise noted above, maps were originally published in Zeiner, D.C., W.F. Laudenslayer, Jr., K.E. Mayer, and M. White, eds. 1988-1990. California's Wildlife. Vol. I-III. California Depart. of Fish and Game, Sacramento, California. Updates are noted in maps that have been added or edited since original publication.

California Wildlife Habitat Relationships System
California Department of Fish and Game
California Interagency Wildlife Task Group

CHIPPING SPARROW

Spizella passerina

Family: EMBERIZIDAE
B489

Order: PASSERIFORMES

Class: AVES

Written by: S. Granholm

Reviewed by: L. Mewaldt, D. Airola

Edited by: R. Duke

DISTRIBUTION, ABUNDANCE, AND SEASONALITY

A common migrant and summer visitor throughout most of California, excluding Central Valley, southern deserts, and alpine areas. Winters less commonly in Central Valley and southern California lowlands. Prefers open wooded habitats with a sparse or low herbaceous layer and few shrubs, if any. Although apparently requires trees for resting and singing, and prefers trees for nesting, often forages in nearby herbaceous and open shrub habitats, including dry margins of wet meadows. Less common in breeding season in southern and interior foothills than in montane habitats, northern coastal ranges, and Great Basin. Fairly common in Imperial and Colorado River valleys in winter, but rare elsewhere. May breed or winter in orchards (Grinnell and Miller 1944, McCaskie et al. 1979, Garrett and Dunn 1981).

SPECIFIC HABITAT REQUIREMENTS

Feeding: Eats mostly insects and spiders in breeding season, and mostly grass and forb seeds remainder of year (Martin et al. 1961). Feeds insects and spiders to young. Gleans from ground and low plants; rarely hawks insects in air. In eastern Texas, perched 30-40 cm (12-16 in) above ground, peered at lower vegetation, and then dropped on insects (Allaire and Fisher 1975).

Cover: Mostly uses trees for nesting, resting, singing, and other cover; also shrubs and ground herbage used (Grinnell and Miller 1944).

Reproduction: In California, usually nests in a conifer, but deciduous tree or shrub also used (Grinnell and Miller 1944). Nest rarely more than 12 m (40 ft) above ground; usually 0.3 to 1.8 m (1-6 ft) or, rarely, on ground (Bent 1968). Nest usually concealed in dense foliage near end of branch.

Water: Drinks water regularly (Smyth and Coulombe 1971), and sometimes bathes in streams (Hering 1948), but can subsist for 3 wk on dry seeds only (Ehrlich et al. 1988).

Pattern: Frequents open woodlands with sparse or low herbaceous cover and a low density of shrubs, if any. Often forages in herbaceous or open shrub habitats nearby.

SPECIES LIFE HISTORY

Activity Patterns: Yearlong, diurnal activity.

Seasonal Movements/Migration: Breeders mostly leave California in September or early October and return in April or May. Some move downslope to winter from Central Valley to southern Mexico, but it is not known what portion of breeding population remains to winter in the state and what portion migrates farther south. Present year-round in parts of southwestern California, but breeding population may be replaced or augmented by a different

wintering population. In portions of winter range not occupied in breeding season, present mostly from September to April.

Home Range: No information found.

Territory: In Michigan, territory varied from 0.4 to 0.6 ha (1 to 1.5 ac), and a territory measured by Sutton (1960) was 0.25 ha (0.65 ac). A pair in Georgia defended 3.1 ha (7.6 ac) during nest-building and incubation, but only 1.1 ha (2.7 ac) while feeding nestlings (Odum and Kuenzler 1955). In Minnesota, territory varied from 0.2 to "about" 0.4 ha (0.5 to 1 ac) (Bent 1968). Average territory in an Arizona pinyon-juniper-ponderosa pine ecotone was 2.7 ha (6.7 ac) (Laudenslayer and Balda 1976).

Reproduction: Breeding season usually begins in April or May. A monogamous, solitary nester; clutch size 3-5, usually 4. May be double-brooded (Terres 1980). Incubation 11-14 days, by female. Altricial young tended by both parents; leave nest at 9-12 days, but do not fly until about 14 days. Apparently breeds first at 1 yr (Harrison 1978).

Niche: Elsewhere in U.S., one of the most common hosts of brown-headed cowbird; more than 600 records reported by Friedmann (1963). Apparently an infrequent host in Sierra Nevada. Only 1 of 64 nests or family groups contained cowbird eggs or young (Rothstein et al. 1980, Verner and Ritter 1983, Airola 1986). May wander or drift upslope in family groups postbreeding. May forage in small flocks in winter with juncos, Brewer's sparrows, and other sparrows (Ehrlich et al. 1988).

REFERENCES

- Airola, D. A. 1986. Brown-headed cowbird parasitism and habitat disturbance in the Sierra Nevada. *J. Wildl. Manag.* 50:571-575.
- Alliaire, P. N., and C. D. Fisher. 1975. Feeding ecology of three resident sympatric sparrows in eastern Texas. *Auk* 92:260-269.
- Bent, A. C. (O. L. Austin, Jr., ed.). 1968. Life histories of North American cardinals, grosbeaks, buntings, towhees, finches, sparrows, and allies. 3 Parts. U.S. Natl. Mus. Bull. 237. 1889pp.
- Ehrlich, P. R., D. S. Dobkin, and D. Wheye. 1988. *The birder's handbook*. Simon and Schuster, New York. 785pp.
- Friedmann, H. 1963. Host relations of the parasitic cowbirds. U.S. Natl. Mus. Bull. 233. 276pp.
- Garrett, K., and J. Dunn. 1981. *Birds of southern California*. Los Angeles Audubon Soc. 408pp.
- Grinnell, J., and A. H. Miller. 1944. *The distribution of the birds of California*. Pac. Coast Avifauna No. 27. 608pp.
- Harrison, C. 1978. *A field guide to the nests, eggs and nestlings of North American birds*. W. Collins Sons and Co., Cleveland, OH. 416pp.
- Hering, L. 1948. Nesting birds of the Black Forest, Colorado. *Condor* 50:49-56.
- Laudenslayer, W. F., Jr., and R. P. Balda. 1976. Breeding bird use of a pinyon-juniper-ponderosa pine ecotone. *Auk* 93:571-586.
- Martin, A. C., H. S. Zim, and A. L. Nelson. 1961. *American wildlife and plants, a guide to wildlife food habits*. Dover Publ., Inc., New York. 500pp.
- McCaskie, G., P. De Benedictis, R. Erickson, and J. Morlan. 1979. *Birds of northern California, an annotated field list*. 2nd ed. Golden Gate Audubon Soc., Berkeley. 84pp.
- Odum, E. P., and E. J. Kuenzler. 1955. Measurement of territory and home range size in birds. *Auk* 72:128-137.
- Rothstein, S. I., J. Verner, and E. Stevens. 1980. Range expansion and diurnal changes in dispersion of the brown-headed cowbird in the Sierra Nevada. *Auk* 97:253-267.
- Smyth, M., and H. M. Coulombe. 1971. Notes on the use of desert springs by birds in California. *Condor* 73:240-243.
- Sutton, G. M. 1960. The nesting fringillids of the Edwin S. George Reserve, southeastern Michigan (VI). *Jack-Pine Warbler* 37:127-151.

- Terres, J. K. 1980. The Audubon Society encyclopedia of North American birds. A. Knopf, New York. 1100pp.
- Verner J., and L. V. Ritter. 1983. Current status of the brown-headed cowbird in the Sierra National Forest. *Auk* 100:355-368.
- Walkinshaw, L. H. 1944. The eastern chipping sparrow in Michigan. *Wilson Bull.* 56:193-205.

B489

Life history accounts for species in the California Wildlife Habitat Relationships (CWHR) System were originally published in: Zeiner, D.C., W.F.Laudenslayer, Jr., K.E. Mayer, and M. White, eds. 1988-1990. California's Wildlife. Vol. I-III. California Department of Fish and Game, Sacramento, California. Updates are noted in accounts that have been added or edited since original publication.